

More integrals of products of Airy functions

Francisco M. Fernández

INIFTA (UNLP, CCT La Plata-CONICET), División Química Teórica, Blvd. 113
S/N, Sucursal 4, Casilla de Correo 16, 1900 La Plata, Argentina

E-mail: fernande@quimica.unlp.edu.ar

Abstract. By means of a modified hypervirial theorem we derive simple expressions for the integrals of products of Airy functions. Present results contain earlier ones as particular cases.

Airy functions appear in many problems of mathematical physics. For that reason some time ago Gordon [1] and Albright [2] derived integrals of products of Airy functions that do not appear in commonly available tables of integrals and functions. However, they are necessary for several problems of physical interest [1, 2].

The purpose of this letter is to propose a simple approach for the derivation of such integrals. It is based on a straightforward modification of the well known hypervirial theorems [3] and may also be useful for other similar problems.

Let $A(x)$ and $B(x)$ be solutions to the eigenvalue equations

$$\begin{aligned}\hat{L}A &= aA, \quad \hat{L}B = bB \\ \hat{L} &= \hat{D}^2 - x, \quad \hat{D} = \frac{d}{dx}\end{aligned}\tag{1}$$

We can easily prove that for any linear operator $\hat{O}$

$$\int A[\hat{L}, \hat{O}]B \, dx = (a - b) \int A\hat{O}B \, dx + W(A, \hat{O}B)\tag{2}$$

where $[\hat{L}, \hat{O}] = \hat{L}\hat{O} - \hat{O}\hat{L}$ is the commutator, $W(u, v) = uv' - u'v$ is the Wronskian and the prime indicates differentiation with respect to x .

We need the following two cases

$$[\hat{L}, f\hat{D}] = f''\hat{D} + 2f'(\hat{L} + x) + f\tag{3}$$

and

$$[\hat{L}, g] = g'' + 2g'\hat{D}\tag{4}$$

where f and g are differentiable functions of x . It follows from equation (2) with $\hat{O} = f\hat{D}$ and $\hat{O} = g$ that

$$\int Af''\hat{D}B \, dx + 2b \int Af'B \, dx + 2 \int Ax f'B \, dx + \int AfB \, dx =\tag{5}$$

$$(a - b) \int Af\hat{D}B \, dx + W(A, fB')\tag{6}$$

and

$$\int Ag' \hat{D}B \, dx = \frac{a-b}{2} \int AgB \, dx - \frac{1}{2} \int Ag''B \, dx + \frac{1}{2} W(A, gB) \quad (7)$$

respectively. Notice that we can use the latter equation twice with $g = f'$ and with $g' = f$ to remove the two integrals in the former equation that contain $\hat{D}B = B'$. Thus we derive the following expression

$$\begin{aligned} \int AfB \, dx + 2 \int Ax f' B \, dx - \frac{1}{2} \int Af''' B \, dx + (a+b) \int Af' B \, dx \\ - \frac{(a-b)^2}{2} \int AFB \, dx = W(A, fB') - \frac{1}{2} W(A, f'B) \\ + \frac{a-b}{2} W(A, FB) \end{aligned} \quad (8)$$

where $F = \int f \, dx$. From this master equation we easily obtain any integral of the form $\int x^n A(x)B(x) \, dx$ such as the particular cases considered by Gordon [1]. Notice that he chose functions of the form $A[\alpha(R + \beta_1)]$ but we can set the scale $\alpha = 1$ without loss of generality so that $\beta_1 = a$ and $\beta_2 = b$ in our notation. For example, if $F = 1$ we obtain his equation (B.10):

$$\int A(x)B(x) \, dx = \frac{1}{b-a} [A'(x)B(x) - A(x)B'(x)] \quad (9)$$

If we choose $a = b = 0$ then the master equation (8) gives us all the integrals of the form $I_n(y_1, y_2) = \int x^n y_1 y_2 \, dx$ considered by Albright [2]. For example, when $f = x^n$ we obtain the recurrence relation

$$\begin{aligned} \int x^n A(x)B(x) \, dx = \frac{n(n-1)(n-2)}{2(2n+1)} \int x^{n-3} A(x)B(x) \, dx + \frac{x^{n+1} A(x)B(x)}{2n+1} \\ - \frac{n(n-1)x^{n-2} A(x)B(x)}{2(2n+1)} + \frac{nx^{n-1} [A(x)B'(x) + A'(x)B(x)]}{2(2n+1)} \\ - \frac{x^n A'(x)B'(x)}{2n+1}, \quad n = 0, 1, \dots \end{aligned} \quad (10)$$

On the other hand, equation (7) enables us to connect the integrals $I_n(y_1, y'_2)$ with the $I_m(y_1, y_2)$ ones. Finally, if we take into account that

$$\int f y'_1 y'_2 \, dx = f y'_1 y_2 - \int f' y'_1 y_2 \, dx - \int x f y_1 y_2 \, dx \quad (11)$$

we easily express the integrals $I_k(y'_1, y'_2)$ in terms of those discussed above.

In our opinion the modified hypervirial theorems provide a most convenient simple and systematic approach to the calculation of integrals of products of Airy functions.

- [1] Gordon R G 1969 *J. Chem. Phys.* **51** 14.
- [2] Albright J R 1977 *J. Phys. A* **10** 485.
- [3] Fernández F M and Castro E A 1987 *Hypervirial Theorems* in Lecture Notes in Chemistry Vol. 43 (Springer-Verlag, Berlin).